

HERBOTIQ



• Category and Product form

Herbal antibiotic activity feed supplement – Powder

• Composition

Mixture of dried herbal extracts

• Mode of action

Herbotiq® can inhibit bacterial virulence factors, inhibit bacterial biofilm formation and prevent bacterial adhesion to host receptors through the bacterial quorum sensing regulation system. As a photosensitizer.

• Dosage and Administration

POULTRY: As a supportive therapy

Regular use: 250 gm / ton of feed.

During mild infections: 500 gm/ton of feed 5 days in month

During disease out breaks: 1kg/ton of feed 7-10 days (or) as advised by a poultry consultant.

Special Note : It should be used mainly as growth promoter and to control mild infections only. During serious outbreaks potential antibiotics should be used.

• Indications

Arrests the growth of bacteria, virus etc present in the gut of the animals.

Reduces the chances of occurrence of microbial infections.

Effectively controls all bacterial infections of poultry like coli bacillosis, fowl typhoid, fowl cholera, coryza, CRD etc.

Helps to reduce the virulence or activity or effectiveness of the virus during out breaks.

Most suitable to eradicate drug resistant bacteria.

• Precautions

For veterinary Use Only.

The product is safe for the intended use

No withdrawal period is necessary

No side effects after administration

No residue in meat

No drug resistance

• Storage

Store in a cool dry place in tightly closed container.

Keep out of reach of children.

Keep away from direct sunlight.

• Packaging

5, 10 and 15 Kg bags

• Shelf Life

2 years in original packaging

