

Composition Each kilogram of powder contains: Zoalene (Dinitolmide) 250g

Target Animal

Broiler, Layer Pullets and Turkey

Dosage and Administration

/ Broilers: Mix 500 g Dinicox® well per ton of feed through the whole cycle.

/ Replacement pullets: Mix 300-500 g Dinicox® well per ton of feed from day old to 16 weeks of age.

/ Breeder: Mix 500 g Dinicox® well per ton of feed, up to 14 weeks of age.

/ Turkey: Mix 500 to 700 g of Dinicox® well per ton of feed from day old to 16 weeks of age.

Note: It is recommended to mix Dinicox® first with 5 times its weight with feed and add the mixture to total feed volume and mix thoroughly again.

Contraindications

Do not use with other coccidiostats

Withdrawal period

/ Meat: Zero Day

/ Egg: Do not use in layers producing eggs for human consumption.

Storage

Keep it in room temperature and away from direct sunlight.

Precautions

/ Keep out of reach of children.

/ Not to be fed to laying birds.

/ Avoid inhaling dust.

/ Avoid contact with eyes and epithelial membrane

Packaging

0.5, 1,5 Kg bag

Environmental Considerations

/ Avoid release to the environment.

/ Dispose in a safe manner in accordance with local regulations.



Indications

/ Broiler Chickens: For prevention and control of coccidiosis

/ Replacement pullet: For development of active immunity to coccidiosis

/ Growing Turkeys: For prevention and control of coccidiosis