

DOXINIQ10%

Antibiotic Solution for oral administration | For veterinary use only

ANTIBIOTICS

Composition Each liter of solution contains: **Doxycycline Hyclate 116 g (Equivalent to 100 g Doxycycline Base)**

Target Animal

Broiler, Breeder and Layer pullet

Indications

To prevent, control and treat diseases caused by microorganisms sensitive to doxycycline including:

- / Chronic Respiratory Disease (CRD) caused by Mycoplasma
- / Joint Infections
- / Pasteurellosis (fowl cholera) caused by Pasteurella multocida
- / Colibacillosis
- / Infectious Coryza caused by Haemophilus
- / Salmonellosis
- / Necrotic Enteritis caused by Clostridia
- / Air Sac Infection
- / Staphylococcus and Streptococcus infections
- / Chlamydia infections
- / Respiratory infections caused by Ornithobacterium rhinotracheale

Dosage and Administration

In Broiler, Breeder and Layer pullet:

/ 0.5-1 ml of Doxiniq®10% per liter of drinking water (equivalent to 10 - 20 mg Doxycycline Base Per each Kg B.W. per day) for 3 to 5 consecutive days.

/ Note: Duration and dosage based on the severity of the disease should be prescribed by a clinician.

Contraindications:

- / Do not administer to laying birds four weeks before the start of lay and during lay.
- / Do not use in animals with hepatic dysfunction.

Packaging

1 liter bottle

Withdrawal period

Poultry meat: 7 days

Drug interactions

Avoid simultaneous use of Doxiniq®10% with Florfenicol, Beta-lactams, Cephalosporins, Aminoglycosides and Cations with 2 and 3 capacities such as Aluminum, Calcium, Magnesium, Zinc and Iron.

Storage

- / Keep it in room temperature (between 15-25 °C) and away from direct sunlight.
- / Keep the lid of container closed.

Precautions

- / Use under Super-vision of a veterinarian.
- / Keep out of reach of children.
- / The shelf life of the medicine after mixing with water is 24 hours.

