

Composition Each liter of solution contains: Norfloxacin 200g

Target Animal

Broiler and Layer Pullet

Indications

Noraxin® 20% is prescribed in poultry for treatment of:

- / Respiratory infections and CRD
- / Gastrointestinal infections and necrotic enteritis
- / Genitourinary infections
- / Mycoplasma infections
- / Septicemia, peritonitis, arthritis
- / Secondary infections caused by viral diseases
- / All other infections caused by gram-negative and gram-positive bacteria sensitive to Norfloxacin such as Salmonella, Fowl Cholera (Pasteurella), Staphylococcus, Streptococcus, Pseudomonas aeruginosa, Campylobacter, Haemophilus, E. coli,

Dosage and Administration

- / Poultry: 1 L Noraxin® 20% / 1500 - 4000 L drinking water (2000 L drinking water on average), during all day, for 3 - 5 consecutive days
- / Duration and dosage should be prescribed by a veterinarian.
- / Medicated drinking water should be used within 24 hours.

Contraindications

None

Withdrawal period

Chicken | Meat: 4 days

Storage

- / Keep it in room temperature and away from direct sunlight.
- / Keep the lid of container closed.

Packaging

1 liter bottle

Precautions

- / Use under supervision of veterinarian.
- / Keep out of reach of children.
- / Shake well before Use.

Environmental Considerations

- / Avoid release to the environment.
- / Dispose in a safe manner in accordance with local regulations.

Incompatibility with other drugs

Do not use simultaneously with tetracyclines, macrolides, lincosamides, chloramphenicol.

