

Composition Each liter of Solution contains: Ofloxacin 50g

Target Animal

Poultry

Dosage and Administration

Mix OFL cure® into the drinking water at 0,1 – 0,2 ml for each kg of bodyweight for 3-5 days.

/ Duration and dosage should be prescribed by a veterinarian.

Contraindications

/ Do not administer to animals with renal and liver dysfunction.

/ Do not administer to laying hens producing eggs for human consumption

Withdrawal period

Meat & offal: 7 days

Incompatibility with other drugs

Do not use simultaneously with tetracyclines, macrolides, lincosamides, chloramphenicol, and non-steroidal anti-inflammatory preparations.

Storage

/ Store in cool and dry place (5 - 25°C).

/ Avoid direct sunlight.

/ Keep the lid of container closed.

/ Use the product up to 28 days after opening the seal of container.

Precautions

/ Keep out of reach of children.


/ Do not administer to laying birds 14 days before the start of lay and during lay.

Packaging

1 liter bottle

Environmental Considerations

Dispose in a safe manner in accordance with local regulations.



Indications

OFL cure® is a broad spectrum antibacterial from second generation of fluoroquinolones. It can be used for treatment of infections caused by gram-positive, gram-negative bacteria, Mycoplasma, and in the case of CRD and CRD complex, Colibacillosis, Coryza, Salmonellosis, Omphalitis and Fowl Cholera.