

# Tiamphenicure 10%

Broad spectrum antibiotic | For Veterinary Use Only

ANTIBIOTICS

**Composition** Each liter of solution contains: **Tiamphenicol 100 g**

## Target Animal

Poultry, Livestock

## Indications

Tiamphenicure 10%® is indicated for the treatment of respiratory infections, Salmonellosis, gastro-enteritis, meningitis and encephalitis caused by micro-organisms susceptible to tiamphenicol in calves and poultry.

## Dosage and Administration

/ General dosage: 30 - 40 mg tiamphenicol / kg b.w. For oral administration via drinking water or milk replacer.

/ Calves: 2 - 3 ml / liter of drinking water, daily, for up to 5 days.

/ Poultry: 1.5 - 2 L / 1000 liter of drinking water, daily, for up to 5 days.

Duration and dosage should be prescribed by a veterinarian.

Medicated drinking water should be used within 24 hours. Medicated milk replacer should be used within 12 hours.

## Contraindications

/ Do not administer to animals hypersensitive to tiamphenicol.

/ Do not administer to animals with an active rumen and intestinal microbial flora.

/ Do not administer to animals with an impaired kidney function.

Do not use in layers producing eggs for human consumption.

## Packaging

1 liter bottle

## Withdrawal period

Meat: 12 days from the last treatment

Milk: 5 days from the last treatment

## Storage

/ Keep it in room temperature and away from direct sunlight.

/ Keep the lid of container closed.

## Precautions

/ Keep out of reach of children.

/ Use in pregnant or lactating animals only after risk/benefit assessment by veterinarian.

/ Reduce dosage in animals with impaired kidney function.

/ Do not use after Exp date.

/ The shelf life, if sealed and subject to storage conditions, is 3 years from the date of manufacture and 28 days if opened.

## Environmental Considerations

/ Avoid release to the environment.

/ Dispose in a safe manner in accordance with local regulations.

