Broad spectrum antibiotic | For Veterinary Use Only

# **Composition Each liter of solution contains: Thiamphenicol 250 g**

### **Target Animal**

Poultry, Turkey

### **Indications**

/ Broiler: treatment of bacterial disease of the respiratory and intestinal tract, in chronic respiratory disease and in fowl coryza, sustained in particular by Gram-positive and Gram-negative microorganisms, ricketsiae, mycoplasms and chlamydiae (mycoplasma spp. Pasturella spp. Staphylococcus spp. Salmonella spp., ornithobacterium rhinotracheale, clostridium perfringens).

/ Turkey: infections sustained by salmonella spp, pasturella multocida, ornithobacterium rhinotracheale, staphylococcus aureus, clostridium perfringens, mycoplasma gallisepticum, mycoplasma synoviae, mycoplasma meleagridis.

### **Contraindications**

/ Do not administer to animals hypersensitive to thiamphenicol.

/ Do not administer to animals with an active rumen and intestinal microbial flora.

/ Do not administer to animals with an impaired kidney function.

/ Do not use in layers producing eggs for human consumption.

# Withdrawal period

Broiler: 4 days Turkey: 6 days

# Storage

/ Keep it in room temperature and away from direct sunlight.

/ Keep the lid of container closed.

# **Incompatibility with other drugs**

Do not combine with penicillin, cephalosporins, amino-glycosides, as it will result in an antagonistic effect.

## **Precautions**

/ Keep out of reach of children.

/ Use in pregnant or lactating animals only after risk/benefit assessment by veterinarian.

/ Reduce dosage in animals with impaired kidney function.

/ Do not use after Exp date.

/ The shelf life, if sealed and subject to storage conditions, is 3 years from the date of manufacture and 28 days if opened.

#### **Environmental Considerations**

/ Avoid release to the environment.

/ Dispose in a safe manner in accordance with local regulations

