

Composition Each 100 grams of powder contains: Tylvalosin 62.5 g

Target Animal

Broiler, Breeder and Layer pullet

Indications

It's used for treatment of respiratory and enteric diseases caused by Tylvalosin sensitive microorganism, mainly Broiler, Breeder and Layer pullet, such as: Mycoplasma pneumonia, Ornithobacteriosis and necrotic enteritis.

Dosage and Administration

In drinking water:

Broiler, Breeder and Layer pullet:

25 mg Tylvalosin for 1 kg of body weight of a bird every twenty - four hours over the course of 3 days.

Note: First dissolve each 100 g sachet in 10 - 15 L of drinking water then add it to the main drinking water tank.

Contraindications

No incompatibility has been found between Tylvacare® and other water-soluble pharmaceuticals and/or feed additives, which are normally used for poultry.

Withdrawal period

Meat: 2 days.

Eggs: zero days.

Storage

/ Keep it in a dry and dark room with temperature 15 °C to 25 °C.

/ Keep out of reach of children.

/ Use by date after initial opening of primary packaging 5 weeks.

Precautions

Once dissolved in water the prepared solution should be used within 24 hours.

It is advised to use fresh solution daily.

Direct contact with eyes, skin and mouth should be avoided and protective equipment should be worn.

Skin that has been in contact with Tylvacare® or feed or drinking water that contains Tylvacare® should be washed.

Environmental Considerations

Dispose empty sachets in a safe manner in accordance with local regulations.

Packaging

100 g Sachet

