

Composition Each Kilogram of product contains: Salinomycin 120g

Target Animal

Poultry

Indications

- / Do not use simultaneously with tiamulin and other coccidiostats.
- / Do not use for breeding poultry, turkeys, ducks, horses and other solid-hoofed animals, ruminants.
- / Do not use for laying hens.
- / Do not leave or dispose of food with the product (Salinomycin) in places available for farm or wildlife animals.

Dosage and Administration

- / Broiler: Add 500 g Uniq Salino® 12% to 1 ton of feed and mix it well.
- / Layer pullets: Add 350-500 g Uniq Salino® 12% to 1 ton of feed and mix it well. Drug must be consumed by week 16th and 7 days before laying process.
- / To obtain a uniform mixture, first mix the required quantity with 20-50 kg of feed and add the mixture to the main feed volume and mix it again.

Contraindications

- horses and other solid-hoofed animals, ruminants.
- / Do not use for laying hens.
- / Do not leave or dispose of food with the product (Salinomycin) in places available for farm or wildlife animals.

Withdrawal period

- / Meat and offal: 5 days

Storage

- / Keep it in room temperature and away from direct sunlight.
- / Keep the container closed.

Precautions

- / For veterinary use only.
- / Keep out of reach of children.

Packaging

1,5,10,15 and 25 Kg bags

