



Disinfectant Solution Based on Aldehydes and Quaternary Ammonium Compounds | For Veterinary Use Only

Composition Each liter of solution contains: **Didecyl Dimethyl Ammonium Chloride 20g | Glutaraldehyde 60g**
Formaldehyde 30g | Benzalkonium Chloride 50g

DISINFECTANT

Target Animal

Livestock and poultry

Dosage and Administration

First prepare diluted solution in accordance with the instruction below:

- / Fungicidal effect: Add 0.5 liters of Virover® into 99.5 liters water.
- / Bactericidal effect: Add 750 ml of Virover® into 99.25 liters water.
- / Virucidal effect: Add 1 liter of Virover® into 99 liters water.
- / Then spray 100 - 200 ml of diluted solution per square meter of surfaces.

Storage

- / Store in a cool place, away from direct sunlight and keep lid of container closed.
- / Should be used in 60 days after opening the lid of container.
- / Should be used in 48 hours after preparing the diluted solution.

Precautions

- / Protect eyes and skin by using suitable clothes, gloves, mask and eyeglasses.
- / Avoid swallowing the solution and inhaling its vapors and spray particles.
- / This product is irritant to skin, eyes and mucous membranes, if contacting wash immediately with plenty of clean water and refer to physician.
- / Keep out of reach of children.

Packaging

5 Liter Gallon

Environmental Considerations

- / This material and its container must be disposed of in a safe way.
- / Empty containers may retain some product residues and hence be potentially hazardous.
- / Dispose of waste to licensed waste disposal site in accordance with the requirements of the local regulations.

