

Animal
Health

"Discover the Unique Cure – Precision in
Every Drop, Every Dose."



export@uniqcure.com

"Antibiotic Solution for Oral Administration in Poultry –
Contains Doxycycline for Effective Treatment."

Doxiniq 10%

www.uniqcure.com

Doxiniq® 10%

Antibiotic Solution for oral administration

For veterinary use only

Composition

Each liter of solution contains:

Doxycycline Hyclate 116 g
(equivalent to 100 g Doxycycline Base)

Target Animal

Broiler, Breeder and Layer pullet

Indication

To prevent, control and treat diseases caused by microorganisms sensitive to doxycycline including:

- Chronic Respiratory Disease (CRD) caused by Mycoplasma
- Joint Infections
- Pasteurellosis (fowl cholera) caused by Pasteurella multocida
- Colibacillosis
- Infectious Coryza caused by Haemophilus
- Salmonellosis
- Necrotic Enteritis caused by Clostridia
- Air Sac Infection
- Staphylococcus and Streptococcus infections
- Chlamydia infections
- Respiratory infections caused by Ornithobacterium rhinotracheale

Dosage and Administration

In Broiler, Breeder and Layer pullet:

0.5 - 1 ml of Doxiniq®10% / liter of drinking water (equivalent to 10-20 mg Doxycycline Base / Kg B.W. / day) for 3 to 5 consecutive days.

Note:

- Medicated water must be used within 12 hours during therapy period.
- Dosage and Duration should be prescribed by a veterinarian.

Contraindication

- Do not administer to laying birds four weeks before the start of lay and during lay.
- Do not use in animals with hepatic dysfunction.
- Do not use for layers producing eggs for human consumption.

Withdrawal period

Meat (Broiler): 5 days

Drug interaction

Avoid simultaneous use of Doxiniq®10% with Penicillins, Cephalosporins, Aminoglycosides and Cations with 2 and 3 valencies such as Aluminum, Calcium, Magnesium, Zinc and Iron.

Storage

- Keep it in room temperature and away from direct sunlight.
- Keep the lid of container closed.

Precaution

- Use under supervision of a veterinarian.
- Keep out of reach of children.
- Do not use after Exp date.
- The shelf life, if sealed and subject to storage conditions, is 2 years after manufacturing date and 28 days if opened.

Packaging

1 liter bottle