

Animal Health

"Discover the Unique Cure – Precision in Every Drop, Every Dose."



export@uniqcure.com

"Antibiotic Solution for Oral Administration in Poultry – Contains Florfenicol for Effective Treatment."

Florniq 10%

www.uniqcure.com

Florniq® 10%
Antibiotic solution for oral administration
For veterinary use only

Composition
Each liter of solution contains:
Florfenicol 100 g

Target animal
Broiler, breeder and layer pullet

Indication
Prevention and treatment of Gastrointestinal and Respiratory infections caused by Florfenicol susceptible microorganisms such as: Actinobacillus spp., Pasteurella spp., Salmonella spp. and Streptococcus spp. in poultry.

Dosage and Administration
1 L of Florniq® 10% / 1000 liter of drinking water (Equivalent to 20 mg of Florfenicol / Kg body weight) daily for 3-5 consecutive days.
Note:

- This product does not need to be diluted in a small amount of water. To prevent active substance precipitation in drinking water, add the calculated amount of Florniq® 10% directly to the drinking water tank.
- Medicated water must be used within 24 hours during therapy period.
- Dosage and duration should be prescribed by a veterinarian.

Contraindication
Do not use for layers producing eggs for human consumption.

Withdrawal Period
Chicken meat: 5 days

Drug interaction
Avoid the simultaneous use of Florfenicol with other antibiotics such as Penicillin, Macrolides and Aminoglycosides.

Storage

- Keep it in room temperature and away from direct sunlight.
- Keep the lid of container closed.

Precaution

- Use under supervision of a veterinarian.
- Keep out of reach of children.
- Do not use after Exp date.
- The shelf life, if sealed and subject to storage conditions, is 2 years after manufacturing date and 28 days if opened.

Packaging
1 liter bottle