

Animal Health

"Discover the Unique Cure – Precision in Every Drop, Every Dose."



export@uniqcure.com

"Antibiotic Solution for Oral Administration in Poultry – Contains Norfloxacin for Effective Treatment."

Noraxin 20%

www.uniqcure.com

Noraxin®20%

Antibiotic solution for oral administration

For Veterinary Use Only

Composition

Each liter of solution contains:

Norfloxacin 200 g

Target Animal

Broiler and Layer Pullet

Indication

- Treatment of Respiratory infections and CRD
- Treatment of Gastrointestinal infections and Necrotic enteritis
- Treatment of Genitourinary infections
- Treatment of Mycoplasma infections
- Septicemia, Peritonitis, Arthritis
- Secondary infections caused by viral diseases
- Other infections caused by gram-negative and gram-positive bacteria sensitive to Norfloxacin such as Salmonella, Fowl Cholera (Pasteurella), Staphylococcus, Streptococcus, Pseudomonas aeruginosa, Campylobacter, Haemophilus, E. coli.

Dosage and Administration

1 L Noraxin® 20% / 1500 - 4000 L of drinking water, (2000 L of drinking water on average), during all day, for 3 - 5 consecutive days

Note:

- Duration and dosage should be prescribed by a veterinarian.
- Medicated water must be used within 24 hours during therapy period.

Contraindication

Do not use for layers producing eggs for human consumption.

Withdrawal time

Chicken meat: 4 days

Drug interaction

Avoid the simultaneous use of Norfloxacin with Tetracyclines, Macrolides, Lincosamides and Chloramphenicol.

Storage

- Keep it in room temperature and away from direct sunlight.
- Keep the lid of container closed.

Precaution

- Use under supervision of a veterinarian.
- Keep out of reach of children.
- Do not use after Exp date.
- The shelf life, if sealed and subject to storage conditions, is 2 years after manufacturing date and 28 days if opened.

Packaging

1 Liter Bottle