

Animal
Health

"Discover the Unique Cure – Precision in
Every Drop, Every Dose."



export@uniqcure.com

"Three-Component Antibiotic (Water-Soluble Powder –
WSP) for Oral Administration in Poultry"

Sulfaves

www.uniqcure.com

Sulfaves®

Water soluble powder antibiotic for oral administration

For Veterinary Use Only

Composition

Each 100 gr of product contains:

- Sodium Sulfamerazine Activity 40 g
- Sodium Sulfamethazine Activity 40 g
- Sodium Sulfaquinoxaline Activity 20 g

Target Animal

Poultry and Turkey

Indication

As an aid to control and treat Coccidiosis, Pasteurellosis (acute fowl cholera) and Salmonellosis in chickens and turkeys caused by pathogens susceptible to sulfamerazine, sulfamethazine and sulfaquinoxaline.

Dosage and Administration

1) Coccidiosis - Chicken:

- As an aid to control and treat coccidiosis caused by E.tenella, E.mivati, E.maxima, E.brunetti, E.acervulina and E.necatrix.

Provide medicated water (0.04% solution) for 2-3 days, then plain water for 3 days, then medicated water (0.025% solution) for 2 days.

- If bloody droppings appear, repeat at 0.025% level for 2 more days.

- Do not change litter during therapy period.

2) Acute Fowl Cholera - Chicken and Turkey:

- As an aid to control and treat acute fowl cholera caused by Pasteurella multocida, Provide medicated water (0.04% solution) for 2-3 days.

- If disease recurs, repeat treatment.

3) Salmonellosis - Chicken and Turkeys:

As an aid to control and treat salmonellosis caused by Salmonella species, Provide medicated water (0.033% solution) for 5 days.

4) Coccidiosis - Turkey:

- As an aid to control and treat coccidiosis caused by E.meleagritidis and E.adenooides, Provide medicated water (0.025% solution) for 2 days, then plain water for 3 days, then medicated water (0.025% solution) for 2 days, then plain water for 3 days, then medicated water (0.025% solution) for 2 days.

- Repeat if necessary.

- Do not change litter during therapy period.

Instruction for preparation of 3Sulfachick® solution:

- 0.04% Solution: Add one 100 gr sachet of 3Sulfachick® to 250 liters drinking water.
- 0.033% Solution: Add one 100 gr sachet of 3Sulfachick® to 300 liters drinking water.
- 0.025% Solution: Add one 100 gr sachet of 3Sulfachick® to 400 liters drinking water.

Note:

- Make fresh solution daily.
- Use only medicated water during treatment unless otherwise directed.
- For control of disease outbreaks, should start 3Sulfachick® treatment as soon as the disease is diagnosed.
- If improvement is not noted in 72 hrs., consult your veterinarian.
- Treated animals must actually consume enough medicated water to provide a therapeutic dose.

Contraindication

Don't use in layers producing eggs for human consumption.

Withdrawal period

Chicken and Turkey meat: 14 days

Drug interaction

None.

Storage

- Keep it in room temperature, dry place and away from direct sunlight.
- Keep the sachet closed.

Precaution

- Use under supervision of veterinarian.
- Keep out of reach of children.
- Do not mix or administer in galvanized containers.

Packaging

100 g sachet