

Animal Health

"Discover the Unique Cure – Precision in Every Drop, Every Dose."



export@uniqcure.com

"Antibiotic Solution for Oral Administration in Poultry – Contains Thiamphenicol for Effective Treatment."

Thiamphenicure 25%

www.uniqcure.com

Thiamphenicure®25%
Antibiotic solution for oral administration
For Veterinary Use Only

Composition

Each liter of solution contains:

Thiamphenicol 250 g

Target Animal Broiler and Turkey

Indication

- Broiler: Treatment of bacterial disease of the respiratory and intestinal tract, Chronic Respiratory Disease, Colibacillosis, Fowl Coryza, treatment of infectious diseases sustained by Gram-positive and Gram-negative microorganisms, Rickettsia, Chlamydia, Mycoplasma spp., Pasteurella spp., Staphylococcus spp., Salmonella spp., Ornithobacterium rhinotracheale and Clostridium perfringens.
- Turkey: Treatment of infectious diseases sustained by Salmonella spp., Pasteurella multocida, Ornithobacterium rhinotracheale, Staphylococcus aureus, Clostridium perfringens, Mycoplasma gallisepticum, Mycoplasma synoviae, Mycoplasma meleagridis.

Dosage and Administration

1.6-2 ml of Thiamphenicure®25% /10 Kg BW (Equivalent to 40-50 mg of thiamphenicol/Kg BW), daily for 3-5 consecutive days in drinking water.

Note:

- The concentration of diluted solution mustn't be less than 800 ml/1000 L D.W. and more than 1250 ml/1000 L D.W.
- This product does not need to be diluted in a small amount of water. To prevent active substance precipitation in drinking water, add the calculated amount of Thiamphenicure®25% directly to the drinking water tank.
- Medicated water must be used within 12 hours during therapy period.
- Duration and dosage should be prescribed by a veterinarian.

Contraindication

- Do not use for animals with an impaired kidney function.
- Do not use in layers producing eggs for human consumption.

Withdrawal Period

Meat and offal

- Broiler: 4 days
- Turkey: 6 days

Drug Interaction

Avoid the simultaneous use of thiamphenicol with Penicillins, Cephalosporines and Aminoglycosides.

Storage

- Keep it in room temperature and away from direct sunlight.
- Keep the lid of container closed.

Precaution

- Use under supervision of a veterinarian.
- Keep out of reach of children.
- Do not use after Exp date.
- The shelf life, if sealed and subject to storage conditions, is 2 years after manufacturing date and 28 days if opened.

Packaging

1 Liter Bottle