

Feed
Additive

"Discover the Unique Cure – Precision in
Every Drop, Every Dose."



export@uniqcure.com

"A powerful blend of activated ascorbic acid and organic acids with antimicrobial properties, designed to support respiratory and gastrointestinal health in poultry."

Ultrax

www.uniqcure.com

Ultrax® Liquid

Mixture of activated Ascorbic acid and Organic acids with antimicrobial effects on respiratory and gastrointestinal system

Feed additive in drinking water and spray

For Veterinary Use Only

Composition

Each liter of solution contains:

- Activated Ascorbic acid 25 g
- Lactic acid 35 g
- Citric acid 30 g

Target animal

Poultry and Livestock

Indication, dosage and administration

1) Prevention and help to treat respiratory diseases caused by fungal, bacterial and viral agents

- 2 ml of Ultrax® Liquid/ 10 L of drinking water for 4-5 consecutive days simultaneous with antibiotic
- 1 ml of Ultrax® Liquid/ 1 L of water per each 250 m², Spray 2-4 times a day, at least for 4-5 consecutive days

2) Prevention and control of post-vaccination reactions

- 1 ml of Ultrax® Liquid/ 10 L of drinking water 12 hrs. after vaccination with Live respiratory vaccines such as Newcastle and Bronchitis, for at least 3 consecutive days
- 1 ml of Ultrax® Liquid/ 1 L of water per each 250 m², Spray 2 times a day 12 hrs. after vaccination with Live vaccines, for at least 3 consecutive days

Note: Do not use Ultrax® Liquid 24 hours before and 12 hours after vaccination.

3) Prevention and help to treat Gastrointestinal diseases

2-3 ml of Ultrax® Liquid/ 10 L of drinking water for 3-7 consecutive days

4) Coadjutant to treatment of mycoplasmosis infected flocks

- 1 ml of Ultrax® Liquid/ 1 L of drinking water, twice a day, each time lasts for 8 hrs. for 3-5 consecutive days, simultaneous with anti-mycoplasmosis medicines
- Also 1 ml of Ultrax® Liquid/ 1 L of water, Spray for 3-5 consecutive days simultaneous with above instruction.

5) Prevention and help to treat Salmonellosis

- Prevention: 1 ml of Ultrax® Liquid/ 10 L of drinking water continuously
- Coadjutant to treatment: 2 ml of Ultrax® Liquid/ 10 L of drinking water for at least 21 consecutive days

Note: Dosage and duration should be prescribed by a veterinarian.

For more information about other indications and dosage refer to the product catalogue.

Contraindication

None.

Withdrawal period

None.

Storage

- Keep it in room temperature and away from direct sunlight.
- Keep the lid of container closed.

Precaution

- Keep out of reach of children.
- Protect eyes and skin by using suitable clothing (gloves, mask and eyeglasses) while using the product.

Packaging

1 L bottle