



UNIQCURE

ADDITIVES & PHARMA

about us.

Since 2018, we have been dedicated to research and development in the production of veterinary drugs and supplements. By 2021, we successfully designed and manufactured 70 products, and by 2024, we expanded our portfolio to 120 products. Looking ahead, our vision is to develop and produce high-quality medicines and supplements for pets, livestock, and aquatic species.

We are now seeking to collaborate with distributors across various countries. If you are interested in learning more about our products and partnership opportunities, please feel free to reach out to us.

visit us.

www.uniqcure.com

contact us.

export@uniqcure.com



BRONCHONIQ®

BRONCHONIQ®
Emulsion Based on High-Purity
Natural Herbal Essential Oils,
Vitamin A & Vitamin C
Full Protection of Poultry
Respiratory System
For Veterinary Use Only

Composition

- Each liter of emulsion contains:
- Eucalyptus Essential Oil
 - Peppermint Essential Oil
 - Menthol
 - Vitamins

Target Animal
Poultry

Indications

- BRONCHONIQ® is indicated for:
- Evacuation of respiratory tract secretions, improved oxygenation, and reduction of suffocation symptoms
 - Use under conditions of high ammonia, hydrogen sulfide, and dust concentration
 - Supportive treatment of upper respiratory tract inflammation
 - Cooling, soothing, and moisturizing respiratory tract during heat stress
 - Reduction of phlegm accumulation and facilitation of its expulsion
 - Cases of anorexia and reduced feed intake
 - Use during recovery period after diseases
 - Control of post-vaccination reactions
 - Strengthening the immune system and repair of damaged respiratory mucosa
 - Acts as a bronchodilator and expectorant

Dosage and Administration

BRONCHONIQ® can be administered via drinking water or by spray.
Prevention & Supportive Treatment of Respiratory Diseases

- Drinking Water:
200 ml BRONCHONIQ® per 1000 liters of drinking water
For 3–5 consecutive days

- Spray Application:
200 ml BRONCHONIQ® per 10 liters of lukewarm water
For 3–5 consecutive days

Control of Post-Vaccination Reactions

- Drinking Water:
200 ml BRONCHONIQ® per 1000 liters of drinking water
Starting 2 days after vaccination, twice daily for 3 consecutive days

Contraindications

Avoid use 1 day before and 2 days after administration of live vaccines.

Withdrawal Period

None.

Drug Interactions

None reported.

Storage Conditions

- Store at room temperature
- Protect from direct sunlight
- Keep the container tightly closed

Precautions


- Keep out of reach of children
- Shake well before use

Environmental Considerations

- Avoid release into the environment
- Dispose of empty containers and waste safely according to local regulations

Packaging

1 Liter Bottle



Improve
oxygenation, and
reduction of
suffocation
symptoms

