



UNIQCURE

ADDITIVES & PHARMA

about us.

Since 2018, we have been dedicated to research and development in the production of veterinary drugs and supplements. By 2021, we successfully designed and manufactured 70 products, and by 2024, we expanded our portfolio to 120 products. Looking ahead, our vision is to develop and produce high-quality medicines and supplements for pets, livestock, and aquatic species.

We are now seeking to collaborate with distributors across various countries. If you are interested in learning more about our products and partnership opportunities, please feel free to reach out to us.

visit us.

www.uniqcure.com

contact us.

export@uniqcure.com



DICLAZUNIQ® 2.5%

Diclazuril – Antiparasitic Oral Solution
For Veterinary Use Only

Composition

Each L contains:

- Diclazuril 25 g (2.5%)
- Excipients q.s.

- Target Animal
- Poultry (Broilers, Layers, Breeders)
 - Livestock (as advised by veterinarian)

- Indication
- DICLAZUNIQ® 2.5% is a highly effective anticoccidial solution indicated for the treatment and prevention of coccidiosis caused by Eimeria species in poultry and livestock, including:
- Eimeria tenella, Eimeria acervulina, Eimeria maxima, Eimeria necatrix, Eimeria brunetti
- It helps to:
- Control acute and subclinical coccidiosis
 - Reduce intestinal damage and bloody diarrhea
 - Improve feed conversion and weight gain
 - Lower mortality and morbidity
 - Enhance overall flock performance

- Dosage and Administration
- Poultry (via drinking water):
- 30-67 ml/ 1000L of Drinking water
- Severe or high-risk infections:
- Treatment may be repeated for 3 consecutive days, if necessary, under veterinary supervision.
- Ensure birds consume only medicated water during treatment.
- Prepare fresh solution daily.
- Livestock:
- 1ml/ 25kg of body weight
 - Treatment may be repeated after 3 weeks, if necessary, under veterinary supervision.

- Note
- Use at early stages of coccidiosis for best results.
 - Maintain strict hygiene and litter management.
 - Suitable for use in shuttle and rotational anticoccidial programs.
 - Can be used safely alongside vitamins and electrolytes.

- Withdrawal Period
- Meat: 1 – 2 days
 - Eggs: Not for human consumption during treatment

- Drug Interaction
- No significant interactions reported at recommended doses
 - Avoid mixing with other anticoccidials unless advised

- Storage
- Store below 25°C
 - Protect from light
 - Do not freeze
 - Keep the bottle tightly closed

- Precaution
- For veterinary use only
 - Follow veterinarian guidance for large-scale use
 - Keep out of reach of children
 - Do not contaminate water sources

Environmental Consideration

Dispose of empty containers safely according to local environmental regulations.

Packaging

1 Liter Bottle



Controls acute and subclinical coccidiosis

