



UNIQCURE

ADDITIVES & PHARMA

about us.

Since 2018, we have been dedicated to research and development in the production of veterinary drugs and supplements. By 2021, we successfully designed and manufactured 70 products, and by 2024, we expanded our portfolio to 120 products. Looking ahead, our vision is to develop and produce high-quality medicines and supplements for pets, livestock, and aquatic species.

We are now seeking to collaborate with distributors across various countries. If you are interested in learning more about our products and partnership opportunities, please feel free to reach out to us.

visit us.

www.uniqcure.com

contact us.

export@uniqcure.com



DINICOX®

DINICOX®
Anticoccidiosis Premix Containing
Zoalene (Dinitolmide)
For Veterinary Use Only

Composition

Composition
Each 1 kg of product contains:
Zoalene (Dinitolmide) 250 g

Target Animals

- Broilers
- Layer Pullets
- Turkeys

Indications

DINICOX® is used for:

- Broilers: Prevention and control of coccidiosis
- Layer pullets: Development of active immunity against coccidiosis
- Growing turkeys: Prevention and control of coccidiosis

Dosage and Administration

- Broilers:
500 g DINICOX® / ton of complete feed, continuously throughout the whole production cycle.
- Layer Pullets:
300–500 g DINICOX® / ton of complete feed, continuously from day-old to 16 weeks of age.
- Growing Turkeys:
500–700 g DINICOX® / ton of complete feed, continuously from day-old to 16 weeks of age.

Mixing Instructions:

It is recommended to premix DINICOX® with 5 times its weight of feed, then add to the total feed volume and mix thoroughly to ensure uniform distribution.

Contraindications

- Do not use in layers producing eggs for human consumption
- Do not use with other anticoccidial drugs

Withdrawal Period

- Meat: None

Storage Conditions

- Store in a dry, cool place (below 25°C)
- Protect from direct sunlight
- Keep the bag tightly closed

Precautions

- Use under the supervision of a veterinarian
- Avoid direct contact with skin and eyes and inhalation of dust
- Contact may cause irritation and sensitivity
- Use protective clothing, gloves, and dust mask
- In case of eye contact, rinse immediately with plenty of water for 15 minutes
- Keep out of reach of children

Packaging

- 25 kg bag



Prevention and control of coccidiosis

