



# UNIQCURE

## ADDITIVES & PHARMA

### about us.

Since 2018, we have been dedicated to research and development in the production of veterinary drugs and supplements. By 2021, we successfully designed and manufactured 70 products, and by 2024, we expanded our portfolio to 120 products. Looking ahead, our vision is to develop and produce high-quality medicines and supplements for pets, livestock, and aquatic species.

We are now seeking to collaborate with distributors across various countries. If you are interested in learning more about our products and partnership opportunities, please feel free to reach out to us.

### visit us.

[www.uniqcure.com](http://www.uniqcure.com)

### contact us.

[export@uniqcure.com](mailto:export@uniqcure.com)





# DOXINIQ® 10%

DOXINIQ® 10%  
Doxycycline Oral Solution  
For Veterinary Use Only

## Composition

Each liter of solution contains:  
Doxycycline 100 g

Target Animals  
Broilers, Breeders, and Layer Pullets

Indications  
DOXINIQ® 10% is used to prevent, control, and treat infections caused by microorganisms sensitive to doxycycline, including:

- Chronic Respiratory Disease (CRD) caused by Mycoplasma
- Joint infections
- Pasteurellosis (Fowl Cholera) caused by Pasteurella multocida
- Colibacillosis
- Infectious Coryza caused by Haemophilus
- Salmonellosis
- Necrotic Enteritis caused by Clostridia
- Air Sac infections
- Staphylococcus and Streptococcus infections
- Chlamydia infections
- Respiratory infections caused by Ornithobacterium rhinotracheale

Dosage and Administration  
Broilers, Breeders & Layer Pullets:  
0.5 – 1 ml of DOXINIQ® 10% per liter of drinking water  
(Equivalent to 10–20 mg Doxycycline Base per kg body weight per day)  
For 3–5 consecutive days

Notes

- Medicated water must be used within 12 hours during the therapy period
- Dosage and duration should be prescribed by a veterinarian

Contraindications

- Do not administer to laying birds four weeks before the start of lay and during lay
- Do not use in animals with hepatic dysfunction
- Do not use in layers producing eggs for human consumption

Withdrawal Period

- Meat (Broilers): 5 days

Drug Interactions  
Avoid simultaneous use of DOXINIQ® 10% with:

- Penicillins, Cephalosporins, and Aminoglycosides
- Divalent and trivalent cations such as Aluminum, Calcium, Magnesium, Zinc, and Iron

Storage Conditions

- Store at room temperature
- Protect from direct sunlight
- Keep the container tightly closed

Precautions

- Use under the supervision of a veterinarian
- Keep out of reach of children
- Do not use after the expiry date
- Shelf life:
  - 2 years when sealed under proper storage conditions
  - 28 days after first opening

Environmental Considerations

- Avoid release into the environment
- Dispose of empty containers and waste safely according to local regulations

Packaging  
1 Liter Bottle



Prevent, control,  
and treat  
infections caused  
by  
microorganisms  
sensitive to  
doxycycline

