

# UNIQCURE ADDITIVES & PHARMA

## about us.

Since 2018, we have been dedicated to research and development in the production of veterinary drugs and supplements. By 2021, we successfully designed and manufactured 70 products, and by 2024, we expanded our portfolio to 120 products.

Looking ahead, our vision is to develop and produce high-quality medicines and supplements for pets, livestock, and aquatic species.

We are now seeking to collaborate with distributors across various countries. If you are interested in learning more about our products and partnership opportunities, please feel free to reach out to us.

**visit us.**

[www.uniqcure.com](http://www.uniqcure.com)

**contact us.**

[export@uniqcure.com](mailto:export@uniqcure.com)



# FOSFUNIQ® 25%

FOSFUNIQ® 25%  
Broad-Spectrum Water-Soluble  
Antibiotic Powder  
For Veterinary Use Only

## Composition

### Composition

Each 1 kg of powder contains:  
Fosfomycin 250 g

#### Target Animals

- Broilers
- Breeders
- Turkeys

#### Indications

FOSFUNIQ® 25% is used as a first-choice antibiotic for the prevention and treatment of the following diseases in broilers, breeders, and turkeys:

- Colibacillosis
- Omphalitis
- Salmonellosis
- Infectious Coryza
- Pasteurellosis
- Staphylococcal arthritis
- Ornithobacteriosis
- Ulcerative and necrotic enteritis
- Staphylococcal and Streptococcal infections

#### Highly sensitive bacteria include:

Escherichia coli, Salmonella, Pasteurella, Citrobacter, Enterococcus, Staphylococcus, Streptococcus, Proteus, Pseudomonas, Clostridium, Vibrio, Aeromonas, Mycobacterium, Haemophilus, Klebsiella, Shigella, Enterobacteriaceae, Serratia, and Ornithobacterium.

#### Dosage and Administration

##### 1) Therapeutic Dose

- 160 mg FOSFUNIQ® 25% / kg BW / day
- Administer in drinking water for 3–5 consecutive days

##### Daily Dose Calculation (g/day):

Average Body Weight (kg) × Number of Birds × 0.16

##### 2) Preventive Dose

- 80 mg FOSFUNIQ® 25% / kg BW / day
- Administer in drinking water for 5–10 consecutive days

##### Daily Dose Calculation (g/day):

Average Body Weight (kg) × Number of Birds × 0.08

#### Method of Adding to Drinking Water

- First dissolve the required amount of FOSFUNIQ® 25% in a few liters of water (18–22°C).
- Add the prepared solution to the main drinking water tank in the morning.
- Ensure the medicated water is fully consumed within 6–8 hours.
- After consumption, provide fresh medicine-free water for the rest of the day.

#### Contraindications

- Do not use in laying birds producing eggs for human consumption.

#### Withdrawal Period

- Poultry meat: 7 days

#### Drug Interactions

- None reported
- Synergistic effects observed with:
  - Erythromycin
  - Tylosin
  - Tylosin
  - Ampicillin
  - Trimethoprim

#### Storage Conditions

- Store at room temperature
- Keep in a dry place
- Protect from direct sunlight
- Keep the sachet tightly closed

#### Precautions

- Use under the supervision of a veterinarian
- Keep out of reach of children

#### Environmental Considerations

- Avoid release into the environment
- Dispose of unused product and empty sachets safely according to local regulations

#### Packaging

- 500 g sachet



## Treatment of bacterial infections

