



# UNIQCURE

## ADDITIVES & PHARMA

### about us.

Since 2018, we have been dedicated to research and development in the production of veterinary drugs and supplements. By 2021, we successfully designed and manufactured 70 products, and by 2024, we expanded our portfolio to 120 products. Looking ahead, our vision is to develop and produce high-quality medicines and supplements for pets, livestock, and aquatic species.

We are now seeking to collaborate with distributors across various countries. If you are interested in learning more about our products and partnership opportunities, please feel free to reach out to us.

### visit us.

[www.uniqcure.com](http://www.uniqcure.com)

### contact us.

[export@uniqcure.com](mailto:export@uniqcure.com)





# HEPANIQ® POWDER

HEPANIQ® POWDER  
Mixture of Herbal Extract & Choline Chloride for Protecting and Improving Poultry Liver Function  
For Veterinary Use Only

## Composition

Each kilogram of powder contains:

- Artichoke Extract
- Choline Chloride

Target Animals  
Broilers, Breeders, and Layers

Indications  
HEPANIQ® Powder is used to:

- Support and protect the liver against toxins such as mycotoxins and certain medicines
- Strengthen and improve liver function
- Stimulate regeneration of hepatic cells
- Support the treatment of fatty liver syndrome in poultry
- Increase bile duct contraction power and bile volume
- Improve digestion and nutrient absorption in the gastrointestinal tract
- Improve feed conversion ratio (FCR) and feed consumption
- Improve egg production in breeders and layers
- Improve hatchability in breeders
- Improve flock uniformity

Dosage and Administration  
General Dose: 50–250 g HEPANIQ® Powder per ton of feed

Broilers

- 200 g/ton: From day 1 to 21
- 100 g/ton: From day 21 to 35
- 50 g/ton: From day 35 to market age

Breeders

- 200 g/ton: From 20 to 40 weeks of age
- 100 g/ton: From 40 to 60 weeks of age
- 50 g/ton: From 60 weeks to end of production

Layers

- 200 g/ton: From 18 to 38 weeks of age
- 100 g/ton: From 38 to 50 weeks of age
- 50 g/ton: From 50 weeks to end of production

Additional Instructions:

- During antibiotic or medicinal treatments, add 50 g extra per ton to the above dosages
- It is recommended to premix HEPANIQ® Powder with 10 times its weight of feed, then blend into total feed thoroughly

Contraindications  
None known.

Withdrawal Period  
None.

Drug Interactions  
None.

Storage Conditions

- Store at room temperature in a dry place
- Protect from direct sunlight
- Keep the sachet tightly closed

Precautions

- Use under the supervision of a veterinarian
- Keep out of reach of children

Environmental Considerations

- Avoid release of product contents into the environment
- Used bags should be disposed of according to local and regional regulations

Packaging  
1 kg , 10kg , 25 kg Bag



Support and protect  
the liver against  
toxins

