

UNIQCURE ADDITIVES & PHARMA

about us.

Since 2018, we have been dedicated to research and development in the production of veterinary drugs and supplements. By 2021, we successfully designed and manufactured 70 products, and by 2024, we expanded our portfolio to 120 products.

Looking ahead, our vision is to develop and produce high-quality medicines and supplements for pets, livestock, and aquatic species.

We are now seeking to collaborate with distributors across various countries. If you are interested in learning more about our products and partnership opportunities, please feel free to reach out to us.

visit us.

www.uniqcure.com

contact us.

export@uniqcure.com



SOLTRINIQ®

SOLTRINIQ®
Antibiotic Oral Solution
For Veterinary Use Only

Composition

Each liter of solution contains:

- Sulfamethoxazole
- Trimethoprim

Target Animal
Broilers

Indications
SOLTRINIQ® is indicated for the treatment and prevention of the following bacterial diseases in broilers caused by microorganisms sensitive to trimethoprim and sulfamethoxazole:

- Colibacillosis caused by *Escherichia coli*
- Infections caused by *Pasteurella multocida*
- Salmonellosis caused by *Salmonella typhimurium* and *Salmonella pullorum*
- Coryza caused by *Avibacterium paragallinarum*

Note: The presence of disease in the flock must be confirmed before treatment.

Dosage and Administration
7.5 mg Trimethoprim and 37.5 mg Sulfamethoxazole per kg body weight per day (Equivalent to 3.75 ml SOLTRINIQ® per 10 kg body weight per day)
Or:
1.5 – 2.5 liters of SOLTRINIQ® per 1000 liters of drinking water
For 3 consecutive days

Notes

- This product does not require pre-dilution in a small amount of water
- To prevent precipitation, add the calculated dose directly to the main water tank
- Prepare fresh solution daily
- Medicated water must be consumed within 24 hours
- During treatment, poultry must not have access to any other water source
- Ensure continuous adequate water availability
- Preferably use low-hardness tap water with pH above 6
- Avoid mixing in galvanized containers
- Dosage and duration must be prescribed by a veterinarian

Contraindications

- Do not use in poultry with severe liver or kidney disease
- Do not use in poultry with impaired hematopoietic systems
- Do not use in layers producing eggs for human consumption
- Avoid simultaneous use with acidic products that reduce stability and solubility

Withdrawal Period

- Poultry meat: 5 days

Drug Interactions

- Do not combine with other veterinary medicinal products without veterinary advice
- Avoid simultaneous use with beta-lactams and amphenicols
- Has synergistic effect with colistin, neomycin, and erythromycin

Storage Conditions

- Store at room temperature
- Protect from direct sunlight
- Keep the container tightly closed
- Shelf life:
 - 2 years when sealed under proper storage conditions
 - 28 days after first opening

Precautions

- Use under the supervision of a veterinarian
- Keep out of reach of children
- Avoid direct contact with eyes, skin, and mucous membranes
- Decreased water consumption may be observed in broilers; use palatability enhancers if necessary to maintain adequate water intake

Environmental Considerations

- Avoid release into the environment
- Dispose of empty containers and waste safely in accordance with local regulations

Packaging
1 Liter Bottle

55

Treatment and
prevention of the
bacterial diseases

