



UNIQCURE

ADDITIVES & PHARMA

about us.

Since 2018, we have been dedicated to research and development in the production of veterinary drugs and supplements. By 2021, we successfully designed and manufactured 70 products, and by 2024, we expanded our portfolio to 120 products. Looking ahead, our vision is to develop and produce high-quality medicines and supplements for pets, livestock, and aquatic species.

We are now seeking to collaborate with distributors across various countries. If you are interested in learning more about our products and partnership opportunities, please feel free to reach out to us.

visit us.

www.uniqcure.com

contact us.

export@uniqcure.com



SULFAVES®

SULFAVES®
Water-Soluble Antibiotic Powder for
Oral Administration
For Veterinary Use Only

Composition

Composition
Each 100 g of product contains:
Sodium Sulfamerazine
Sodium Sulfamethazine
Sodium Sulfaquinoxaline

Target Animals

- Poultry
- Turkeys

Indications

SULFAVES® is indicated as an aid in the control and treatment of the following diseases in chickens and turkeys caused by pathogens susceptible to sulfamerazine, sulfamethazine, and sulfaquinoxaline:

- Coccidiosis
- Pasteurellosis (Acute Fowl Cholera)
- Salmonellosis

Dosage and Administration

1) Coccidiosis – Chickens

As an aid to control and treat coccidiosis caused by E. tenella, E. mivati, E. maxima, E. brunetti, E. acervulina, and E. necatrix:

- Provide medicated water (0.04% solution) for 2–3 days
- Then plain water for 3 days
- Then medicated water (0.025% solution) for 2 days
- If bloody droppings appear, repeat 0.025% solution for 2 more days
- Do not change litter during the treatment period

2) Pasteurellosis (Acute Fowl Cholera) – Chickens & Turkeys

As an aid to control and treat Pasteurella multocida infection:

- Provide medicated water (0.04% solution) for 2–3 days
- If disease recurs, repeat the treatment

3) Salmonellosis – Chickens & Turkeys

As an aid to control and treat Salmonella infections:

- Provide medicated water (0.033% solution) for 5 days

4) Coccidiosis – Turkeys

As an aid to control and treat infections caused by E. meleagrimitis and E. adenoeides:

- Provide medicated water (0.025% solution) for 2 days
- Then plain water for 3 days
- Then medicated water (0.025% solution) for 2 days
- Then plain water for 3 days
- Then medicated water (0.025% solution) for 2 days
- Repeat if necessary
- Do not change litter during the therapy period

Instructions for Preparation of SULFAVES® Solution

- 0.04% Solution:
Add 1 sachet (100 g) to 250 liters of drinking water
- 0.033% Solution:
Add 1 sachet (100 g) to 300 liters of drinking water
- 0.025% Solution:
Add 1 sachet (100 g) to 400 liters of drinking water

Notes

- Prepare a fresh solution daily
- Use only medicated water during treatment unless otherwise directed
- For outbreaks, start SULFAVES® immediately after diagnosis
- If no improvement within 72 hours, consult a veterinarian
- Treated animals must consume sufficient medicated water to receive a therapeutic dose

Contraindications

- Do not use in laying birds producing eggs for human consumption

Withdrawal Period

- Chicken and Turkey meat: 14 days

Drug Interactions

- None known

Storage Conditions

- Store in a dry place at room temperature
- Protect from direct sunlight
- Keep the sachet tightly closed

Precautions

- Use under the supervision of a veterinarian
- Keep out of reach of children
- Do not mix or administer in galvanized containers

Environmental Considerations

- Avoid release into the environment
- Dispose of unused product and empty sachets safely according to local regulations

Packaging

- 100 g sachet



For Treatment of Coccidiosis Pasteurellosis Salmonellosis

