



UNIQCURE

ADDITIVES & PHARMA

about us.

Since 2018, we have been dedicated to research and development in the production of veterinary drugs and supplements. By 2021, we successfully designed and manufactured 70 products, and by 2024, we expanded our portfolio to 120 products. Looking ahead, our vision is to develop and produce high-quality medicines and supplements for pets, livestock, and aquatic species.

We are now seeking to collaborate with distributors across various countries. If you are interested in learning more about our products and partnership opportunities, please feel free to reach out to us.

visit us.

www.uniqcure.com

contact us.

export@uniqcure.com



THIAMPHENICURE® 25%

THIAMPHENICURE® 25%
Thiamphenicol Oral Solution
For Veterinary Use Only

Composition

Each liter of solution contains:
Thiamphenicol 250 g

Target Animals
Broilers and Turkeys

Indications

Broilers:

Treatment of bacterial diseases of the respiratory and intestinal tract, including:

- Chronic Respiratory Disease (CRD)
- Colibacillosis
- Fowl Coryza
- Infections caused by Gram-positive and Gram-negative microorganisms
- Infections caused by Rickettsia, Chlamydia, Mycoplasma spp., Pasteurella spp., Staphylococcus spp., Salmonella spp., Ornithobacterium rhinotracheale and Clostridium perfringens

Turkeys:

Treatment of infectious diseases caused by:

- Salmonella spp.
- Pasteurella multocida
- Ornithobacterium rhinotracheale
- Staphylococcus aureus
- Clostridium perfringens
- Mycoplasma gallisepticum
- Mycoplasma synoviae
- Mycoplasma meleagridis

Dosage and Administration

1.6 – 2 ml of THIAMPHENICURE® 25% per 10 kg body weight daily
For 3–5 consecutive days via drinking water.

Notes

- The concentration of diluted solution must not be less than 800 ml/1000 L and not more than 1250 ml/1000 L of drinking water
- This product does not require pre-dilution in a small amount of water
- Medicated water must be used within 12 hours during the therapy period
- Dosage and duration should be prescribed by a veterinarian

Contraindications

- Do not use in animals with impaired kidney function
- Do not use in layers producing eggs for human consumption

Withdrawal Period (Meat & Offal)

- Broilers: 4 days
- Turkeys: 6 days

Drug Interactions

Avoid simultaneous use with:

- Penicillins
- Cephalosporins
- Aminoglycosides

Storage Conditions

- Store at room temperature
- Protect from direct sunlight
- Keep the container tightly closed

Precautions


- Use under the supervision of a veterinarian
- Keep out of reach of children
- Do not use after the expiry date
- Shelf life:
 - 2 years when sealed under proper storage
 - 28 days after first opening

Environmental Considerations

- Avoid release into the environment
- Dispose of empty containers and waste safely according to local regulations

Packaging

1 Liter Bottle



Treatment of bacterial diseases

