



UNIQCURE

ADDITIVES & PHARMA

about us.

Since 2018, we have been dedicated to research and development in the production of veterinary drugs and supplements. By 2021, we successfully designed and manufactured 70 products, and by 2024, we expanded our portfolio to 120 products. Looking ahead, our vision is to develop and produce high-quality medicines and supplements for pets, livestock, and aquatic species.

We are now seeking to collaborate with distributors across various countries. If you are interested in learning more about our products and partnership opportunities, please feel free to reach out to us.

visit us.

www.uniqcure.com

contact us.

export@uniqcure.com



TYLVACARE®

TYLVACARE®
Water-Soluble Antibiotic Powder for
Oral Administration
For Veterinary Use Only

Composition

Composition
Each 100 g of powder contains:
Tylvalosin Tartrate 62.5 g

Target Animals

- Poultry: Broilers, Breeders, Layers
- Turkeys

Indications

TYLVACARE® is indicated for the prevention and treatment of infections caused by:

- Mycoplasma spp.
- Clostridial infections
- Ornithobacterium rhinotracheale (ORT)

Dosage and Administration (summarized)

Mycoplasma (Prevention & Treatment)

- 40 mg/kg BW/day
 - o Broilers: Days 1–3
 - o Breeders: Days 1–7(≈ 1 sachet / 2500 kg BW/day)
- 24 mg/kg BW/day
 - o Broilers: Days 20–22
 - o Breeders & Layers: Periodic use & during stress(≈ 1 sachet / 4200 kg BW/day)

Clostridial Infections

- 40 mg/kg BW/day for 3–4 days

Ornithobacteriosis (ORT)

- 40 mg/kg BW/day for 5 days

Instructions for Use

- Use only as medicated drinking water during treatment.
- No other water source must be available.
- First dissolve the required amount of TYLVACARE® in 10–15 liters of water (15–25°C), then add to the main water tank.
- Prepare fresh solution daily only for one-day use.

Contraindications

- None reported

Withdrawal Period

- Meat: 2 days
- Eggs: None (Use in laying poultry is permitted)

Storage Conditions

- Store at below 25°C (ideal: 15–25°C)
- Store in a dry place, protected from direct sunlight
- Use within 5 weeks after opening the sachet
- Use within 24 hours after adding to drinking water
- Keep the sachet tightly closed

Precautions

- Use under supervision of a veterinarian
- Keep out of reach of children
- Avoid direct contact with eyes, skin, and mouth
- Wear protective equipment
- Wash exposed skin, eyes, or mucous membranes thoroughly with plenty of water

Packaging

- 100 g Sachet



Treatment of Mycoplasma spp. Clostridial infections Ornithobacterium rhinotracheale (ORT)

