



UNIQCURE

ADDITIVES & PHARMA

about us.

Since 2018, we have been dedicated to research and development in the production of veterinary drugs and supplements. By 2021, we successfully designed and manufactured 70 products, and by 2024, we expanded our portfolio to 120 products. Looking ahead, our vision is to develop and produce high-quality medicines and supplements for pets, livestock, and aquatic species.

We are now seeking to collaborate with distributors across various countries. If you are interested in learning more about our products and partnership opportunities, please feel free to reach out to us.

visit us.

www.uniqcure.com

contact us.

export@uniqcure.com



ULTRAX® LIQUID

ULTRAX® LIQUID
Mixture of Activated Ascorbic Acid
& Organic Acids with Antimicrobial
Effects
Feed Additive for Drinking Water &
Spray
For Veterinary Use Only

Composition

Each liter of solution contains:

- Activated Ascorbic Acid
- Lactic Acid
- Citric Acid

Target Animal
Poultry

Indications, Dosage & Administration

1. Prevention and Supportive Treatment of Respiratory Diseases

Caused by fungal, bacterial and viral agents:

• Drinking Water:

2 ml ULTRAX® Liquid per 10 liters of drinking water for 4–5 consecutive days, simultaneously with antibiotics

• Spray Application:

1 ml ULTRAX® Liquid per 1 liter of water per each 250 m²,

Spray 2–4 times daily for at least 4–5 consecutive days

2. Prevention and Control of Post-Vaccination Reactions

• Drinking Water:

1 ml ULTRAX® Liquid per 10 liters of drinking water, starting 12 hours after vaccination for at least 3 consecutive days

• Spray Application:

1 ml ULTRAX® Liquid per 1 liter of water per 250 m²,

Spray twice daily, starting 12 hours after vaccination for at least 3 consecutive days

Note: Do NOT use ULTRAX® Liquid 24 hours before and 12 hours after vaccination.

3. Prevention and Supportive Treatment of Gastro-Intestinal Diseases

• 2–3 ml ULTRAX® Liquid per 10 liters of drinking water for 3–7 consecutive days

4. Coadjutant in the Treatment of Mycoplasmosis-Infected Flocks

• Drinking Water:

1 ml ULTRAX® Liquid per 1 liter of drinking water, twice daily, each administration lasting 8 hours, for 3–5 consecutive days, simultaneously with anti-mycoplasmosis medicines

• Spray:

1 ml ULTRAX® Liquid per 1 liter of water, spray for 3–5 consecutive days alongside drinking water administration

5. Prevention & Supportive Treatment of Salmonellosis

• Prevention:

1 ml ULTRAX® Liquid per 10 liters of drinking water, continuously

• Coadjutant to Treatment:

2 ml ULTRAX® Liquid per 10 liters of drinking water for at least 21 consecutive days

Important Note:

Dosage and duration should be prescribed by a veterinarian.

Contraindications

None known.

Withdrawal Period

None.

Drug Interactions

None reported.

Storage Conditions

- Store at room temperature
- Protect from direct sunlight
- Keep the container tightly closed

Precautions

- Keep out of reach of children
- Protect eyes and skin using suitable protective equipment

Environmental Considerations

- Avoid release into the environment
- Dispose of waste and empty containers safely according to local regulations

Packaging

1 Liter Bottle



With Antimicrobial Effects

